

**ASSOCIATION OF PHYSICAL ACTIVITY LEVEL AND PHYSICAL FITNESS
WITH THE COGNITIVE FUNCTION OF PATIENTS IN HEMODIALYSIS**

**ASSOCIAÇÃO ENTRE O NÍVEL DE ATIVIDADE FÍSICA E A APTIDÃO FÍSICA
COM A FUNÇÃO COGNITIVA DE PACIENTES EM HEMODIÁLISE**

Recebido em: 11/11/2025

Aceito em: 17/04/2026

Publicado em: 28/06/2026

Juliedy Waldow Kupske¹ 

Universidade Federal do Rio Grande do Sul

Thaís Severo Dutra² 

Universidade Cruz Alta

Universidade Regional do Noroeste do Estado do Rio Grande do Sul

Universidade Regional Integrada do Alto Uruguai e das Missões

Cleide Dejaira Martins Vieira³ 

Universidade Cruz Alta

Universidade Regional do Noroeste do Estado do Rio Grande do Sul

Universidade Regional Integrada do Alto Uruguai e das Missões

Moane Marchesan Krug⁴ 

Universidade Regional do Noroeste Estado do Rio Grande do Sul

Paulo Ricardo Moreira⁵ 

Universidade de Cruz Alta

Rodrigo de Rosso Krug⁶ 

Universidade de Cruz Alta

¹ Aluna do Programa de Pós-graduação stricto sensu em título de doutorado em ciências do movimento humano da Universidade federal do Rio Grande do Sul. Porto Alegre, RS, Brasil. E-mail: juliedykupske@hotmail.com

² Aluna do Programa de Pós-graduação stricto sensu em título de doutorado em atenção em integral a saúde da Universidade Cruz Alta em parceria com a Universidade noroeste do Rio Grande do Sul e Universidade Regional Integrada do Alto Uruguai e das Missões. Cruz alta, RS, Brasil. E-mail: thaís.severo@hotmail.com

³ Aluna do Programa de Pós-graduação stricto sensu em título de doutorado em atenção em integral a saúde da Universidade Cruz Alta em parceria com a Universidade noroeste do Rio Grande do Sul e Universidade Regional Integrada do Alto Uruguai e das Missões. Cruz alta, RS, Brasil. E-mail: cleidedmvieira@gmail.com

⁴ Doutora em Educação Física pela Universidade Federal de Santa Catarina (UFSC). Professora do departamento de Humanidades e Educação, Universidade Regional do Noroeste do Rio Grande do Sul (UNIJUI). Santa Rosa, RS, Brasil. E-mail: Moane.krug@unijui.edu.br

⁵ Doutor em Medicina (Nefrologia) pela Universidade Federal do Rio Grande do Sul. Professor do departamento de Saúde e Ciências Agrárias, Universidade de Cruz Alta (UNICRUZ). Cruz Alta, RS, Brasil. Email: Pmoreira@unicruz.edu.br

⁶ Doutor em Ciências Médicas (UFSC). Professor do departamento de Saúde e Ciências Agrárias, Universidade de Cruz Alta (UNICRUZ). Cruz Alta, RS, Brasil. Email: Rkrug@unicruz.edu.br

Abstract: This study aimed to correlate cognitive function with the level of physical activity and physical fitness over a three-year period in patients undergoing hemodialysis and to compare physical activity levels and physical fitness over this time between patients with and without cognitive problems. To this end, a panel study was conducted, including patients evaluated between 2018 and 2020, using the MMSE, physical-functional record, forearm flexion test, sit-to-stand test, handgrip strength test, and a physical activity monitor. The results showed that cognitive function, assessed in different years of the research, correlated positively with physical fitness and physical activity level: in 2018, with handgrip strength ($r = 0.313$; $p = 0.019$); in 2019, with vigorous cardiorespiratory activity ($r = 0.288$; $p = 0.038$), with handgrip strength ($r = 0.235$; $p = 0.033$) and with physical activity level on hemodialysis days ($r = 0.359$; $p = 0.010$) and on non-treatment days ($r = 0.314$; $p = 0.026$); and in 2020, with physical activity level on non-hemodialysis days ($r = 0.387$; $p = 0.014$). It is concluded that patients with higher physical activity levels and better physical fitness had better cognitive function.

Keywords: Chronic Renal Failure; Musculoskeletal and Neural Physiological Phenomena; Physical Functional Performance; Motor Activity; Cognition.

Resumo: Este estudo teve como objetivo correlacionar a função cognitiva com o nível de atividade física e a aptidão física ao longo do período de três anos de pacientes submetidos à Hemodiálise e comparar o NAF e a aptidão física ao longo desse tempo dos pacientes com e sem problemas cognitivos. Para tanto, foi realizado um estudo de painel, incluindo os pacientes avaliados entre 2018 e 2020, por meio do MEEM, prontuário físico-funcional, teste de flexão de antebraço, sentar e levantar, força de preensão manual e pelo monitor de atividade física. Os resultados mostraram que a função cognitiva avaliada nos diferentes anos da pesquisa correlacionou-se positivamente com a aptidão física e com o NAF: em 2018, com a FPM ($r = 0,313$; $p = 0,019$); em 2019, com a aptidão cardiorrespiratória ($r = 0,288$; $p = 0,038$), com a FPM ($r = 235$; $p = 0,033$) e com o NAF em dia de HD ($r = 359$; $p = 0,010$) e em dia sem tratamento ($r = 314$; $p = 0,026$); e em 2020, com o NAF em dia sem HD ($r = 387$; $p = 0,014$). Conclui-se que pacientes com maior NAF e melhor aptidão física apresentaram melhor função cognitiva.

Palavras-chave: Insuficiência Renal Crônica; Fenômenos Fisiológicos Musculoesqueléticos e Neurais; Desempenho Físico Funcional; Atividade Motora; Cognição.

INTRODUCTION

Chronic kidney disease (CKD) is globally considered a common disorder and characterized as a public health problem (Hu; Coresh, 2017). It is currently defined by abnormalities of renal structure or function, assessed and categorized by glomerular filtration rate (GFR), albuminuria, and duration of kidney injury (Glassock; Warnock; Delanaye, 2017).

Its global prevalence is estimated between 5% and 10%, which reveals the dimension of this pathology. The increase is driven mainly by Diabetes Mellitus prevalence, systemic arterial hypertension and obesity, and the aging process (Lv; Zhang, 2019), which are the main risk factors.

In the most advanced stage of CKD, known as chronic renal failure (CRF) - in which the GFR is $< 15 \text{ mL/min/1.73 m}^2$ - hemodialysis (HD) is the predominant method of renal clearance, adopted for 92% of patients on renal replacement therapy (Neves *et al.*, 2020).

Although efficient to maintain the balance of the body, HD treatment involves a monotonous routine with several restrictions, causing psychological impairments, physical and functional deficiencies, and social limitations (Bernardo *et al.*, 2019). In addition, the CKD phenotype is associated with several harmful outcomes to the health of patients, such as

depression (Khan *et al.*, 2019), frailty (Perez *et al.*, 2019), poor quality of life (Marinho *et al.*, 2017), and higher mortality (Teixeira *et al.*, 2015).

Another variable influenced by pathology and treatment is cognitive function. Studies show that the reported prevalence of cognitive impairment in this population ranged from 6.6% to 51% (San *et al.*, 2017). Patients at all stages of CKD are at higher risk of developing cognitive impairment, especially those with $\text{eGFR} < 30 \text{ mL/min/1.73 m}^2$ (Murray *et al.*, 2016), regardless of age-related changes, and tend to worsen with the beginning of dialysis (Tamura *et al.*, 2017).

The mechanisms involved in this picture are not yet established, but some factors such as neuronal damage induced by uremic toxins, ischemic cerebrovascular lesions, oxidative stress, chronic inflammation, anemia, hyperhomocysteinemia, and endothelial dysfunction may be the most significant aggravating factors (Matta *et al.*, 2014).

Considering the effects of hemodialysis treatment and the high rates of cognitive impairment, supervised physical exercise is an essential tool in maintaining the patients' health on HD, influencing several variables (Lacerda *et al.*, 2018). It can improve Kt/V (a method that assesses hemodialysis adequacy/quality), exercise capacity, depression, and life quality, as well as reducing blood pressure without increasing the incidence of adverse events (Pu *et al.*, 2019).

A systematic review that sought to identify factors correlated with cognitive impairment in patients with CKD on HD pointed out that the risk of cognitive impairment can be significantly higher in elderly, women, with stroke, with difficulties in activities of daily living, with minor hemoglobin concentrations, with higher levels of pain, with sleeping difficulties, and with depression (Oh; Mo; Seo, 2019).

The cross-sectional study by Fukushima *et al.* (2019). compared physically active and inactive patients and revealed that the mean total score on the cognitive test was lower than recommended and that those on HD physically active obtained a significant score in the domain of verbal fluency. These results suggest that a higher physical activity level can contribute to a higher cognitive function score, being a potential therapeutic alternative.

On the other hand, cohort epidemiological studies in this population are still scarce, and few have sought to analyze the cognitive function and its associations. In this sense, the present study aimed to correlate cognitive function with the physical activity level (PAL) and physical fitness of patients undergoing HD over three years and to compare the PAL and physical fitness of individuals on HD with and no cognitive problems over the same period.

METHODS

This quantitative panel-type study was carried out at the nephrology clinic of the Hospital São Vicente de Paulo, a reference center in the northwest region of Rio Grande do Sul, located in Cruz Alta, between 2018 and 2020. The population was constituted of 91 patients with CRF on HD from the Renal Clinic of that hospital who met the inclusion criterion, which was to be on hemodialysis for more than three months in 2018.

To calculate the baseline sample size (2018), the formula for calculating prevalence through a simple casual sample was used, plus 20% predicted losses and 15% for association studies (parameters: size of renal clinic population = 91 patients; 95% confidence level; and sampling error = 1 to 5 percentage points), resulting in a sample size of 73 people.

Therefore, all patients were invited to participate in the study. Of these, 30 refused, totaling 61 patients. The refusals are mainly because people report tiredness in all collection attempts and/or that the surveys are very long and exhaustive.

All received guidance about the study and signed the Free and Informed Consent Form (FICF) before being included in the research. The patients underwent cognitive function assessment using the Mini-Mental State Exam (MMSE). In Brazil, this questionnaire was translated and validated by Bertolucci Brucki, Campacci, and Juliano (1994) and consists of 30 questions about temporal and spatial orientation, fixation memory, evocation, attention, calculation, and language (Bertolucci *et al.*, 1994). Illiterate individuals with a score below 20 points and those with formal education with less than 24 points were considered with a probable cognitive deficit (Almeida, 1998).

The functional physical record was also used to collect age information (years) and HD duration (months). To assess the strength and endurance of upper limbs, the forearm flexion test was used: the participants sit on a chair with the back straight and feet on the floor, hold the dumbbell (men: 4 kg; women: 2 kg) with the preferred hand and perform as many push-up reps as they can for 30 seconds; the total is counted by the evaluator and is classified according to the person's gender and age (Rikli; Jones, 2008).

In the case of the lower limbs, the sit-to-stand test was chosen: the patient remains seated in a chair 45 cm high with the back straight, feet flat on the floor and shoulder-width apart; gets up and sits down for 30 seconds, and the maximum number of repetitions is recorded, classified according to gender and age (Rikli; Jones, 2008).

To measure cardiorespiratory capacity, the six-minute walk test was used, which assesses the maximum distance walked during this time, allowing the participant to dictate the

pace of steps and stop walking in case of limiting symptoms (Rikli; Jones, 2008). Patients were classified according to gender and age, as suggested by the authors.

The handgrip test (dynamometry) was applied to measure handgrip strength. Three attempts were made – with a recovery period of one minute between them – and the longest one was used (Bohannon, 2008).

For measuring the physical activity level through the number of daily steps, the OMRON activity monitor model HJA-310° (pedometer) was used. To be considered active, the patients had to perform 4,600 steps or more daily (Tudor-locke *et al.*, 2011). The equipment was delivered to them on the day of treatment before it started and asked to return it in the following HD session, before the start of the session, accounting for two days of use. Thus, people were evaluated on two moments: on a treatment day (first 24 hours) and a non-treatment day (last 24 hours); they were instructed to wear the device in their pocket attached to their clothes and record the complications in a form.

Data collection took place in June 2018 and 2019, and in December 2020, due to the health protocols for coping with Covid-19. First, data normality was tested using the Kolmogorov–Smirnov test, mean and median analysis, and histogram. For characterizing the study sample and the outcome prevalence, descriptive statistics were performed, using absolute and relative frequencies (for qualitative variables) and measures of central tendency and dispersion (for quantitative variables), with the respective confidence intervals of 95% (95%CI).

The correlation of the outcome with the PAL and physical fitness variables was tested individually in each year, based on Spearman's correlation; on the other hand, the comparison of PAL and physical fitness over three years of HD patients with and without cognitive problems was tested using the student's t-test for independent samples or the Mann-Whitney U test, adopting a significance level of 5%.

The study was approved under the consolidated opinion of the Research Ethics Committee No. 4.171.948, CAAE No. 33286120.1.0000.5350. It followed the recommendations of Resolution 466/2012 of the National Health Council, which indicates the procedures for researching with human beings (Brasil, 2012).

RESULTS

Table 01 shows that the mean age of the patients surveyed was 57.65 ± 57.22 years, and the treatment time, 58.72 ± 15.63 months. It can also be seen that the mean cognitive function was 21.62 ± 5.02 (in 2018), 23.15 ± 4.92 (2019), and 24.65 ± 5.13 (2020).

Table 01 – Sociodemographic and health characteristics of patients with CRF on HD. Cruz Alta, Rio Grande do Sul, Brazil (2018, 2019, 2020).

Variables	Mean	Standard deviation
Age in years (n = 61)	57.65	± 57.22
Hemodialysis time in months (n = 61)	58.72	± 15.63
Cognitive function in 2018 (n = 61)	21.62	± 5.02
Cognitive function in 2019 (n = 58)	23.15	± 4.92
Cognitive function in 2020 (n = 40)	24.65	± 5.13

Source: prepared by the authors.

The correlations between cognitive function with PAL and physical fitness can be seen in Table 02. According to these results, it is possible to identify that cognitive function in HD patients was positively and significantly correlated with HGS ($r = 313$, $p = 019$) in the 2018 analysis; with cardiorespiratory fitness ($r = 288$, $p = 038$), handgrip strength ($r = 235$, $p = 033$), physical activity level assessed by the number of steps on hemodialysis day ($r = 359$, $p = 010$), and level of physical activity assessed by the number of steps on the day contrary to hemodialysis ($r = 314$, $p = 026$) in 2019; and with the physical activity level assessed by the number of steps on the day contrary to that of hemodialysis ($r = 387$, $p = 014$) in 2020. These correlations occurred with moderate strength (greater than 0.3) (Leotti *et al.*, 2020).

Table 02 – Correlation between cognitive function with the level of physical activity and physical fitness of patients with CRF on HD over three years. Cruz Alta, Rio Grande do Sul, Brazil (2018, 2019, 2020) (n = 61).

Variables	CF 2018 M $\pm$ SD	R / p	CF 2019 M $\pm$ SD	R / p	CF 2020 M $\pm$ SD	R / p
CRFit	368.2 $\pm$ 166.6	0.240 0.166	367.1 $\pm$ 196.9	0.288 0.038*	390.4 $\pm$ 219.9	0.290 0.069
MRLl	12.5 $\pm$ 6.6	0.139 0.315	13.5 $\pm$ 7.8	0.177 0.201	11.9 $\pm$ 4.8	0.212 0.189
ULMRS	16.1 $\pm$ 5.7	0.186 0.186	16.7 $\pm$ 6.4	0.254 0.069	15.9 $\pm$ 6.3	0.226 0.160
HGS	27.8 $\pm$ 9.7	0.313 0.019*	27.3 $\pm$ 10.0	0.235 0.033*	26.9 $\pm$ 10.4	0.276 0.084
PAL HD	2875.4 $\pm$ 2996.3	0.070 0.620	2644.1 $\pm$ 2571.3	0.359 0.010*	2509.7 $\pm$ 3092.7	0.216 0.181
PAL NHD	3417.8 $\pm$ 3770.9	0.165 0.242	2763.4 $\pm$ 2439.7	0.314 0.026*	1800.4 $\pm$ 2094.1	0.387 0.014*

Source: prepared by the authors.

Notes: CF = cognitive function (Mini-Mental State Exam score); CRFit = cardiorespiratory fitness (distance in meters); MRLL= muscle resistance located in the lower limbs (number of repetitions in 30 seconds); ULMRS = upper limb muscle resistance strength (number of repetitions in 30 seconds); HGS = handgrip strength (KgF); PAL HD = physical activity level assessed by the number of steps on hemodialysis day; PAL NHD = physical activity level assessed by the number of steps on the day contrary to hemodialysis; M = mean; SD = standard deviation; $p \leq 0.05$ for Spearman correlation.

Based on the data presented in Tables 03, 04, and 05, it is noted that HD patients without a probable significant deficit had higher PAL and physical fitness values in all years of analysis compared to those with a probable cognitive deficit. In 2018, this difference was significant in the HGS and NAF HD variables; in 2019, this occurred in ACR and PAL NHD, and in 2020, in CRFit, HGS, PAL HD, and PAL NHD.

Table 03 – Physical activity level and physical fitness of HD patients with and without cognitive problems. Cruz Alta, Rio Grande do Sul, Brazil (2018) (n = 61).

Variables	Cognitive function		
	No probable cognitive deficit (n = 22)	With probable cognitive deficit (n = 18)	p
	M±SD	M±SD	
CRFit	397.09±180.62	340.90±154.53	0.233
MRLL	12.91±7.51	12.23±6.11	0.713
ULMRS	16.78±5.42	15.62±6.09	0.477
HGS	32.43±8.80	24.76±9.19	0.003*
PAL HD	3245.2±3025.4	2625.2±3029.4	0.465**
PAL NHD	4521.4±5003.3	2432.4±1732.0	0.039**

Source: prepared by the authors.

Notes: CRFit = cardiorespiratory fitness (distance in meters); MRLL= muscle resistance located in the lower limbs (number of repetitions in 30 seconds); ULMRS = upper limb muscle resistance strength (number of repetitions in 30 seconds); HGS = handgrip strength (KgF); PAL HD = physical activity level assessed by the number of steps on hemodialysis day; PAL NHD = physical activity level assessed by the number of steps on the day contrary to hemodialysis; M = mean; SD = standard deviation; $p \leq 0.05$ for Spearman correlation; * $p \leq 0.05$ for Student's t-test for independent samples; ** Mann-Whitney U Test was applied.

Table 04 – Physical activity level and physical fitness of HD patients with and without cognitive problems. Cruz Alta, Rio Grande do Sul, Brazil (2019) (n = 58).

Variables	Cognitive function		
	No probable cognitive deficit (n = 23)	With probable cognitive deficit (n = 17)	p
	M±SD (n = 23)	M±SD (n = 17)	
CRFit	426.52±202.94	315.80±182.37	0.048*
MRLL	14.92±8.20	11.92±6.51	0.150
ULMRS	17.38±7.48	16.24±5.38	0.535
HGS	30.04±11.03	25.21±8.93	0.082
PAL HD	3065.40±1763.56	2307.78±3285.47	0.319**
PAL NHD	3633.44±2869.83	1875.30±1596.21	0.013**

Source: prepared by the authors.

Notes: CRFit = cardiorespiratory fitness (distance in meters); MRLL= muscle resistance located in the lower limbs (number of repetitions in 30 seconds); ULMRS = upper limb muscle resistance strength (number of repetitions in 30 seconds); HGS = handgrip strength (KgF); PAL HD = physical activity level assessed by the

number of steps on hemodialysis day; PAL NHD = physical activity level assessed by the number of steps on the day contrary to hemodialysis; M = mean; SD = standard deviation; $p \leq 0.05$ for Spearman correlation; * $p \leq 0.05$ for Student's t-test for independent samples; ** Mann-Whitney U Test was applied.

Table 05 – Physical activity level and physical fitness of HD patients with and without cognitive problems. Cruz Alta, Rio Grande do Sul, Brazil (2020) (n = 40).

Variables	Cognitive function		
	No probable cognitive deficit (n = 27)	With probable cognitive deficit (n = 13)	p
	M±SD	M±SD	
CRFit	455.30±203.71	255.85±195.31	0.006*
MRLL	12.78±4.78	10.08±4.46	0.096
ULMRS	17.15±5.44	13.23±7.21	0.063
HGS	29.00±11.17	22.54±7.24	0.035*
PAL HD	2896.22±3223.91	1707.08±2744.40	0.026**
PAL NHD	2398.74±2314.24	557.77±412.81	0.007*

Source: prepared by the authors.

Notes: CRFit = cardiorespiratory fitness (distance in meters); MRLL= muscle resistance located in the lower limbs (number of repetitions in 30 seconds); ULMRS = upper limb muscle resistance strength (number of repetitions in 30 seconds); HGS = handgrip strength (KgF); PAL HD = physical activity level assessed by the number of steps on hemodialysis day; PAL NHD = physical activity level assessed by the number of steps on the day contrary to hemodialysis; M = mean; SD = standard deviation; $p \leq 0.05$ for Spearman correlation; * $p \leq 0.05$ for Student's t-test for independent samples; ** Mann-Whitney U Test was applied.

DISCUSSION

One of the aims of this study was to correlate cognitive function with PAL and physical fitness of HD patients over three years. The other was to compare the PAL and physical fitness of HD patients with and without cognitive problems in the same period. In this sense, the results showed that those with the best cognitive function were those with higher PAL and better physical fitness and that HD patients without a probable significant deficit had higher PAL and physical fitness values in all years of analysis. In 2018, this difference was significant in the variables handgrip strength (HGS) and physical activity level assessed by the number of steps a day contrary to hemodialysis (PAL NHD); in 2019, in the variables cardiorespiratory fitness and physical activity level assessed by the number of steps on the day contrary to hemodialysis (PAL NHD); and in 2020, in the variables cardiorespiratory fitness, HGS, PAL HD, and PAL NHD.

Similar findings have already been found in other international studies (Fukushima *et al.*, 2019; Manfredin *et al.*, 2017; Stringuetta-Belik *et al.*, 2019; Stringuetta-Belik *et al.*, 2012; Martins *et al.*, 2011). However, none analyzed data longitudinally in a panel format, highlighting their importance in this article. It is known that the prevalence of cognitive impairment is high in this population (Chaiben *et al.*, 2019) and that hemodialysis treatment

contributes to the worsening of this clinical condition (Erken *et al.*, 2019). Studies indicate that cognitive deficit is related to age, years of education, and time on hemodialysis (Gesualdo *et al.*, 2017) and that patients with such impairment are at increased risk of poor health outcomes (Kaltsatou *et al.*, 2015).

As it is a systemic disease, that is, one that affects multiple systems, the correlations found between PAL, physical fitness, and cognitive function can be attributed to several physiological mechanisms arising from physical exercise that interfere with cognitive function. Some pathways are related to immune improvement and inflammation (Müller-Ortiz *et al.*, 2019); activation of NRF2, which acts in neuroprotection against oxidative stressors and mitochondrial toxins (Abreu *et al.*, 2017); improvement in endothelial function, vascular mechanism that affects both the brain and the kidneys (Silva *et al.*, 2019); significant improvement in Kt/V, serum creatinine, serum urea, serum potassium, and phosphorus (Paluchamy; Vaidyanathan, 2018), biochemical markers that influence the clinical picture of the pathology.

It has been pointed out that systematic physical training seems to improve cognitive function in patients with CRF, inducing positive changes in brain metabolism and favoring better scores in their cognitive function (Kaltsatou *et al.*, 2015). Avoiding cognitive impairment through physical training protocols in the routine clinical treatment of HD patients can be a cost-effective alternative, reducing non-adherence to medication, increasing quality of life, and even survival (Schneider *et al.*, 2015).

In this sense, physical activity is potentially related to the physical fitness of hemodialysis patients. It was verified in the present research an improvement in the variable's cardiorespiratory fitness, muscle resistance located in the lower limbs, HGS, which is also correlated with the improvement in cognitive function.

HGS can be considered a protective factor against cognitive decline (Fritz; McCarthy; Adamo, 2019). A cohort study showed that each 1 lb reduction in HGS was associated with a 1.4% increase in the risk of developing neurocognitive pathologies over time (Buchman; Boyle; Wilson; Tang; Bennett, 2007). A stronger handgrip strength may reflect the integrity of the neuromuscular system, resistance to oxidative stress, and inflammation (Weaver *et al.*, 2002). Another work justified this association due to the shared pathology related to hormonal levels (testosterone) and inflammatory biomarkers (IL-6) (Fritz; McCarthy; Adamo, 2019), factors that influence the preservation of cognitive function.

There are explicit links between the motor and cognitive systems (Mcgrath *et al.*, 2019; Bohannon, 2019). The learning of motor skills and motor performance depends on the activity of the frontal and parietal regions of the brain, and the interconnection between these regions is related to motor performance. Thus, improving muscle strength and endurance, that is, physical fitness, reduces the likelihood of developing a cognitive deficit.

The systematic review by Sexton *et al.* (2016), which sought to elucidate the relationship between physical fitness and cognitive function in the elderly, demonstrated that higher levels of physical fitness have beneficial effects on this function. According to the authors, higher levels of physical fitness were associated with larger volumes of gray matter in the brain (GM) and reduced amount or severity of GM lesions or improved measures of GM microstructure.

In the research by Pentikäinen *et al.* (2019), the relationship between cognitive function was evaluated through an extensive battery of neuropsychological tests and cardiorespiratory fitness of 421 individuals with a mean age of 69 years, using peak oxygen consumption (VO₂peak, L/min) for 24 months. Over two years of follow-up, this function decreased from 0.15 to 0.14 points on average, and cardiorespiratory fitness was associated with executive functions and processing speed and also with general cognitive function (Pentikäinen *et al.*, 2019).

Erickson, Leckie, and Weinstein (2014) showed a correlation between physical activity, physical fitness, and cognitive function in adults, elderly, long-lived, and elderly people with different pathologies. Higher levels of physical activity and cardiorespiratory fitness are routinely associated with a larger volume of gray matter in the prefrontal cortex and hippocampus and less consistently in other regions; this suggests that the prefrontal cortex and hippocampus remain flexible at the end of life and that moderate-intensity exercises for six months to a year are sufficient to change the size of these areas (Erickson; Leckie; Weinstein, 2014).

However, it is noteworthy that patients on hemodialysis tend to be less physically active due to the phenotype of the disease. A study using accelerometers identified that people on HD are more sedentary (47.4%) when compared to healthy individuals (10.5%), especially on dialysis days (reduction of 2,645 steps on average) (Gomes *et al.*, 2015).

The daily life of patients on HD requires changes in eating habits, water restrictions, restrictions on travel, work, and activities of daily living, due to the weakness imposed by the

treatment (Araújo *et al.*, 2016). Thus, professionals involved in the health care of chronic renal patients should always be encouraging the regular practice of physical activities.

This research has some limitations, such as the size of the population evaluated, which did not reach the sample size calculation, and sample losses to follow-up. On the other hand, the fact that there is no study in the literature that monitors and assesses the natural history of a condition, such as the correlation and comparison of cognitive function, PAL, and physical fitness over the period of three years of patients undergoing HD, as proposed here.

CONCLUSIONS

It is concluded that the cognitive function of the participants in this study was correlated with the activity level and physical fitness. Patients on HD without probable cognitive deficit had higher values in PAL and physical fitness (HGS, localized muscle resistance of the lower limbs, and cardiorespiratory fitness) in all years of analysis compared to those with probable cognitive deficit. The results indicate physical activity as an effective alternative in improving the physical fitness and cognitive function of these people.

Although this study provides important evidence to health professionals, especially in encouraging patients to participate in training protocols with intradialytic physical exercise, it is suggested that further investigations be carried out using accelerometers or that assess the direct effect of physical exercise practice (such as intervention studies) to reinforce the findings of this work. It is important to highlight that physical activities in environments outside renal therapy units, such as commuting, leisure, and work, and the reduction of sedentary behavior are measures that should be encouraged.

REFERENCES

- ABREU, C. C. *et al.* Does resistance exercise performed during dialysis modulate Nrf2 and NF- κ B in patients with chronic kidney disease? **Life Sciences**, [S. l.], v. 188, p. 192-197, 2017. Disponível em: <https://doi.org/10.1016/j.lfs.2017.09.007>. Acesso em: 8 jun. 2026.
- ARAÚJO, J. B. D. *et al.* Cotidiano de pacientes renais crônicos submetidos à hemodiálise: expectativas, modificações e relações sociais. **Revista de Pesquisa: Cuidado é Fundamental Online**, [S. l.], p. 4996-5001, 2016.
- BERNARDO, M. F. *et al.* Estado nutricional e qualidade de vida de pacientes em hemodiálise. **Medicina (Ribeirão Preto)**, Ribeirão Preto, v. 52, n. 2, p. 128-135, 2019. Disponível em: <https://doi.org/10.11606/issn.2176-7262.v52i2p128-135>. Acesso em: 8 jun. 2026.

BERTOLUCCI, P. H. *et al.* O mini-exame do estado mental em uma população geral: impacto da escolaridade. **Arquivos de Neuro-Psiquiatria**, São Paulo, v. 52, p. 1-7, 1994. Disponível em: <https://doi.org/10.1590/S0004-282X1994000100001>. Acesso em: 8 jun. 2026.

BOHANNON, R. W. Grip strength: an indispensable biomarker for older adults. **Clinical Interventions in Aging**, [S. l.], p. 1681-1691, 2019. Disponível em: <https://doi.org/10.2147/CIA.S194543>. Acesso em: 8 jun. 2026.

BRASIL. Ministério da Saúde. Conselho Nacional de Saúde. Resolução nº 466, de 12 de dezembro de 2012: diretrizes e normas regulamentadoras de pesquisas envolvendo seres humanos. **Diário Oficial da União**, Brasília, DF, v. 150, n. 112, seção 1, p. 59-62, 2013.

BUCHMAN, A. S. *et al.* Frailty is associated with incident Alzheimer's disease and cognitive decline in the elderly. **Psychosomatic Medicine**, [S. l.], v. 69, n. 5, p. 483-489, 2007. Disponível em: <https://doi.org/10.1097/psy.0b013e318068de1d>. Acesso em: 8 jun. 2026.

CHAIBEN, V. B. O. *et al.* Cognition and renal function: findings from a Brazilian population. **Brazilian Journal of Nephrology**, São Paulo, v. 41, p. 200-207, 2018. Disponível em: <https://doi.org/10.1590/2175-8239-JBN-2018-0067>. Acesso em: 8 jun. 2026.

ERICKSON, K. I.; LECKIE, R. L.; WEINSTEIN, A. M. Physical activity, fitness, and gray matter volume. **Neurobiology of Aging**, [S. l.], v. 35, supl., p. S20-S28, 2014. Disponível em: <https://doi.org/10.1016/j.neurobiolaging.2014.03.034>. Acesso em: 8 jun. 2026.

ERKEN, E. *et al.* Impaired cognition in hemodialysis patients: the Montreal Cognitive Assessment (MoCA) and important clues for testing. **Clinical Nephrology**, [S. l.], v. 91, n. 5, p. 275, 2019.

FERRARI, F. *et al.* Exercício físico em indivíduos em hemodiálise: benefícios e melhores indicações – revisão sistemática. **Revista Pesquisa em Fisioterapia**, [S. l.], p. 404-419, 2018. Disponível em: <https://doi.org/10.17267/2238-2704rpf.v8i3.1933>. Acesso em: 8 jun. 2026.

FRITZ, N. E.; MCCARTHY, C. J.; ADAMO, D. E. Handgrip strength as a means of monitoring progression of cognitive decline – a scoping review. **Ageing Research Reviews**, [S. l.], v. 35, p. 112-123, 2017. Disponível em: <https://doi.org/10.1016/j.arr.2017.01.004>. Acesso em: 8 jun. 2026.

FUKUSHIMA, R. L. M. *et al.* Cognitive abilities and physical activity in chronic kidney disease patients undergoing hemodialysis. **Dementia & Neuropsychologia**, São Paulo, v. 13, p. 329-334, 2019. Disponível em: <https://doi.org/10.1590/1980-57642018dn13-030010>. Acesso em: 8 jun. 2026.

GESUALDO, G. D. *et al.* Cognitive impairment of patients with chronic renal disease on hemodialysis and its relationship with sociodemographic and clinical characteristics. **Dementia & Neuropsychologia**, São Paulo, v. 11, p. 221-226, 2017. Disponível em: <https://doi.org/10.1590/1980-57642016dn11-030003>. Acesso em: 8 jun. 2026.

GLASSOCK, R. J.; WARNOCK, D. G.; DELANAYE, P. The global burden of chronic kidney disease: estimates, variability and pitfalls. **Nature Reviews Nephrology**, [S. l.], v. 13,

n. 2, p. 104-114, 2017. Disponível em: <https://doi.org/10.1038/nrneph.2016.163>. Acesso em: 8 jun. 2026.

GOMES, E. P. *et al.* Physical activity in hemodialysis patients measured by triaxial accelerometer. **BioMed Research International**, [S. l.], v. 2015, p. 1-7, 2015. Disponível em: <https://doi.org/10.1155/2015/645645>. Acesso em: 8 jun. 2026.

HU, J. R.; CORESH, J. The public health dimension of chronic kidney disease: what we have learnt over the past decade. **Nephrology Dialysis Transplantation**, [S. l.], v. 32, supl. 2, p. ii113-ii120, 2017. Disponível em: <https://doi.org/10.1093/ndt/gfw416>. Acesso em: 8 jun. 2026.

KALTSATOU, A. *et al.* Cognitive function and exercise training for chronic renal disease patients: a literature review. **Journal of Bodywork and Movement Therapies**, [S. l.], v. 19, n. 3, p. 509-515, 2015. Disponível em: <https://doi.org/10.1016/j.jbmt.2015.04.006>. Acesso em: 8 jun. 2026.

KHAN, A. *et al.* Prevalence and predictors of depression among hemodialysis patients: a prospective follow-up study. **BMC Public Health**, [S. l.], v. 19, n. 1, p. 1-13, 2019. Disponível em: <https://doi.org/10.1186/s12889-019-6796-z>. Acesso em: 8 jun. 2026.

LEOTTI, V. B. *et al.* Modelagem estatística: perguntas que você sempre quis fazer, mas nunca teve coragem. **Clinical and Biomedical Research**, Porto Alegre, v. 39, n. 4, 2019. Disponível em: <https://doi.org/10.22491/2357-9730.98944.98944>. Acesso em: 8 jun. 2026.

LV, J. C.; ZHANG, L. X. Prevalence and disease burden of chronic kidney disease. In: **Renal Fibrosis: mechanisms and therapies**. Singapore: Springer, 2019. p. 3-15. Disponível em: https://doi.org/10.1007/978-981-13-8871-2_1. Acesso em: 8 jun. 2026.

MANFREDINI, F. *et al.* Exercise in patients on dialysis: a multicenter, randomized clinical trial. **Journal of the American Society of Nephrology**, [S. l.], v. 28, n. 4, p. 1259, 2017. Disponível em: <https://doi.org/10.1681/ASN.2016030378>. Acesso em: 8 jun. 2026.

MARINHO, C. L. A. *et al.* **Quality of life of chronic renal patients undergoing hemodialysis**. [S. l.: s. n.], 2017.

MARTINS, C. T. B. *et al.* Comparison of cognitive function between patients on chronic hemodialysis who carry out assisted physical activity and inactive ones. **Brazilian Journal of Nephrology**, São Paulo, v. 33, p. 27-30, 2011. Disponível em: <https://doi.org/10.1590/S0101-28002011000100003>. Acesso em: 8 jun. 2026.

MATTA, S. *et al.* Cognitive alterations in chronic kidney disease: an update. **Jornal Brasileiro de Nefrologia**, São Paulo, v. 36, n. 2, p. 241-245, 2014.

MCGRATH, R. *et al.* Handgrip strength is associated with poorer cognitive functioning in aging Americans. **Journal of Alzheimer's Disease**, [S. l.], v. 70, n. 4, p. 1187-1196, 2019. Disponível em: <https://doi.org/10.3233/JAD-190042>. Acesso em: 8 jun. 2026.

MÜLLER-ORTIZ, H. *et al.* Exercise training in advanced chronic kidney disease. **Revista Médica de Chile**, Santiago, v. 147, n. 11, p. 1443-1448, 2019. Disponível em: <https://doi.org/10.4067/s0034-98872019001101443>. Acesso em: 8 jun. 2026.

MURRAY, A. M. *et al.* The brain in kidney disease (BRINK) cohort study: design and baseline cognitive function. **American Journal of Kidney Diseases**, [S. l.], v. 67, n. 4, p. 593-600, 2016. Disponível em: <https://doi.org/10.1053/j.ajkd.2015.11.008>. Acesso em: 8 jun. 2026.

NEVES, P. D. M. D. M. *et al.* Censo Brasileiro de Diálise: análise de dados da década 2009-2018. **Brazilian Journal of Nephrology**, São Paulo, v. 42, p. 191-200, 2020. Disponível em: <https://doi.org/10.1590/2175-8239-JBN-2019-0234>. Acesso em: 8 jun. 2026.

OH, H.; MO, J.; SEO, W. Correlates of cognitive impairment in patients with chronic kidney failure on haemodialysis: systematic review and meta-analysis. **Journal of Advanced Nursing**, [S. l.], v. 75, n. 5, p. 962-978, 2019. Disponível em: <https://doi.org/10.1111/jan.13907>. Acesso em: 8 jun. 2026.

PALUCHAMY, T.; VAIDYANATHAN, R. Effectiveness of intradialytic exercise on dialysis adequacy, physiological parameters, biochemical markers and quality of life – a pilot study. **Saudi Journal of Kidney Diseases and Transplantation**, Riade, v. 29, n. 4, p. 902-910, 2018. Disponível em: <https://doi.org/10.4103/1319-2442.239661>. Acesso em: 8 jun. 2026.

PENTIKÄINEN, H. *et al.* Cardiorespiratory fitness and cognition: longitudinal associations in the FINGER study. **Journal of Alzheimer's Disease**, [S. l.], v. 68, n. 3, p. 961-968, 2019. Disponível em: <https://doi.org/10.3233/JAD-180897>. Acesso em: 8 jun. 2026.

PEREZ, S. I. A. *et al.* Frailty, health-related quality of life, cognition, depression, vitamin D and health-care utilization in an ambulatory adult population with type 1 or type 2 diabetes mellitus and chronic kidney disease: a cross-sectional analysis. **Canadian Journal of Diabetes**, [S. l.], v. 43, n. 2, p. 90-97, 2019. Disponível em: <https://doi.org/10.1016/j.jcjd.2018.06.001>. Acesso em: 8 jun. 2026.

PU, J. *et al.* Efficacy and safety of intradialytic exercise in haemodialysis patients: a systematic review and meta-analysis. **BMJ Open**, [S. l.], v. 9, n. 1, e020633, 2019. Disponível em: <http://dx.doi.org/10.1136/bmjopen-2017-020633>. Acesso em: 8 jun. 2026.

SAN, A. *et al.* Screening of cognitive impairment in the dialysis population: a scoping review. **Dementia and Geriatric Cognitive Disorders**, [S. l.], v. 44, n. 3-4, p. 182-195, 2017. Disponível em: <https://doi.org/10.1159/000479679>. Acesso em: 8 jun. 2026.

SCHNEIDER, S. M. *et al.* Cognitive function in patients with chronic kidney disease: challenges in neuropsychological assessments. **Seminars in Nephrology**, [S. l.], v. 35, n. 4, p. 304-310, 2015. Disponível em: <https://doi.org/10.1016/j.semnephrol.2015.06.002>. Acesso em: 8 jun. 2026.

SEXTON, C. E. *et al.* A systematic review of MRI studies examining the relationship between physical fitness and activity and the white matter of the ageing brain. **NeuroImage**, [S. l.], v. 131, p. 81-90, 2016. Disponível em: <https://doi.org/10.1016/j.neuroimage.2015.09.071>. Acesso em: 8 jun. 2026.

SILVA, V. R. *et al.* Aerobic exercise training and nontraditional cardiovascular risk factors in hemodialysis patients: results from a prospective randomized trial. **Cardiorenal Medicine**, [S. l.], v. 9, n. 6, p. 391-399, 2019. Disponível em: <https://doi.org/10.1159/000501589>. Acesso em: 8 jun. 2026.

STRINGUETTA-BELIK, F. *et al.* Influence of intradialytic aerobic training in cerebral blood flow and cognitive function in patients with chronic kidney disease: a pilot randomized controlled trial. **Nephron**, [S. l.], v. 140, n. 1, p. 9-17, 2018. Disponível em: <https://doi.org/10.1159/000490005>. Acesso em: 8 jun. 2026.

STRINGUETTA-BELIK, F. *et al.* Maior nível de atividade física associa-se a melhor função cognitiva em renais crônicos em hemodiálise. **Brazilian Journal of Nephrology**, São Paulo, v. 34, p. 378-386, 2012. Disponível em: <https://doi.org/10.5935/0101-2800.20120028>. Acesso em: 8 jun. 2026.

TAMURA, M. K. *et al.* Loss of executive function after dialysis initiation in adults with chronic kidney disease. **Kidney International**, [S. l.], v. 91, n. 4, p. 948-953, 2017. Disponível em: <https://doi.org/10.1016/j.kint.2016.11.015>. Acesso em: 8 jun. 2026.

TEIXEIRA, F. I. R. *et al.* Survival of hemodialysis patients at a university hospital. **Brazilian Journal of Nephrology**, São Paulo, v. 37, p. 64-71, 2015. Disponível em: <https://doi.org/10.5935/0101-2800.20150010>. Acesso em: 8 jun. 2026.

TUDOR-LOCKE, C. *et al.* How many steps/day are enough? For older adults and special populations. **International Journal of Behavioral Nutrition and Physical Activity**, [S. l.], v. 8, n. 1, p. 1-19, 2011. Disponível em: <https://doi.org/10.1186/1479-5868-8-80>. Acesso em: 8 jun. 2026.

WEAVER, J. D. *et al.* Interleukin-6 and risk of cognitive decline: MacArthur studies of successful aging. **Neurology**, [S. l.], v. 59, n. 3, p. 371-378, 2002. Disponível em: <https://doi.org/10.1212/WNL.59.3.371>. Acesso em: 8 jun. 2026.